



SRW

PTO/SB/21 (09-06)

Approved for use through 03/31/2007. OMB 0651-0031
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

TRANSMITTAL FORM

(to be used for all correspondence after initial filing)

Total Number of Pages in This Submission

Application Number	10/562,243
Filing Date	December 22, 2005
First Named Inventor	Garcia-Rodenas et al.
Art Unit	Unknown
Examiner Name	Unknown
Attorney Docket Number	112701-694

ENCLOSURES (Check all that apply)

<input type="checkbox"/> Fee Transmittal Form <input type="checkbox"/> Fee Attached <input type="checkbox"/> Amendment/Reply <input type="checkbox"/> After Final <input type="checkbox"/> Affidavits/declaration(s) <input type="checkbox"/> Extension of Time Request <input type="checkbox"/> Express Abandonment Request <input type="checkbox"/> Information Disclosure Statement <input checked="" type="checkbox"/> Certified Copy of Priority Document(s) <input type="checkbox"/> Reply to Missing Parts/ Incomplete Application <input type="checkbox"/> Reply to Missing Parts under 37 CFR 1.52 or 1.53	<input type="checkbox"/> Drawing(s) <input type="checkbox"/> Licensing-related Papers <input type="checkbox"/> Petition <input type="checkbox"/> Petition to Convert to a Provisional Application <input type="checkbox"/> Power of Attorney, Revocation Change of Correspondence Address <input type="checkbox"/> Terminal Disclaimer <input type="checkbox"/> Request for Refund <input type="checkbox"/> CD, Number of CD(s) _____ <input type="checkbox"/> Landscape Table on CD	<input type="checkbox"/> After Allowance Communication to TC <input type="checkbox"/> Appeal Communication to Board of Appeals and Interferences <input type="checkbox"/> Appeal Communication to TC (Appeal Notice, Brief, Reply Brief) <input type="checkbox"/> Proprietary Information <input type="checkbox"/> Status Letter <input type="checkbox"/> Other Enclosure(s) (please identify below):
Remarks Certified Copy of EP 03014040.4 Return Receipt Postcard		

SIGNATURE OF APPLICANT, ATTORNEY, OR AGENT

Firm Name	Bell, Boyd & Lloyd LLC		
Signature			
Printed name	Robert M. Barrett		
Date	December 11, 2006	Reg. No.	30,142

CERTIFICATE OF TRANSMISSION/MAILING

I hereby certify that this correspondence is being facsimile transmitted to the USPTO or deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450 on the date shown below:

Signature			
Typed or printed name	Heather Foster	Date	December 11, 2006

This collection of information is required by 37 CFR 1.5. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.11 and 1.14. This collection is estimated to 2 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

This Page Blank (uspto)



THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicants: Garcia-Rodenas et al.
Appl. No.: 10/562,243
Filed: December 22, 2005
Conf. No.: 6067
Title: NUTRITIONAL FORMULA FOR OPTIMAL GUT BARRIER
Art Unit: Unknown
Examiner: Unknown
Docket No.: 112701-694

Mail Stop
Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

SUBMISSION OF CERTIFIED COPY OF PRIORITY DOCUMENT

Applicants are respectfully enclosing the certified copy of the priority document for which priority is claimed for the above-identified application under 35 U.S.C. §119. Specifically, the document enclosed is:

<u>Document No.</u>	<u>Country</u>	<u>Date</u>
03014040.4	Europe	June 23, 2003

The Commissioner is hereby authorized to charge deposit account 02-1818 for any fees which are due and owing.

Respectfully submitted,

BELL, BOYD & LLOYD LLC

BY _____

Robert M. Barrett
Reg. No. 30,142
Customer No.: 29157

Dated: December 11, 2006

This Page Blank (uspto)



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03014040.4

**CERTIFIED COPY OF
PRIORITY DOCUMENT**

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

This Page Blank (uspto)



Anmeldung Nr:
Application no.: 03014040.4
Demande no:

Anmeldetag:
Date of filing: 23.06.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

NESTEC S.A.
Avenue Nestlé 55
CH-1800 Vevey
SUISSE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Nutritional formula for optimal gut barrier function

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A23L1/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

This Page Blank (uspto)

Nutritional formula for optimal gut barrier function

Field of the invention

5 The present invention pertains to a composition for inducing a pattern of gut barrier maturation similar to that observed with breast feeding and able to improve gut barrier maturation, e.g. during neonatal stress. In particular, the present invention relates to an infant formula containing a combination of specific ingredients designed to provide a synergistic effect all along
10 gastrointestinal tract and barrier function.

Background of the invention

During the postnatal development, the newborn intestine experiences a process of maturation that ends by the establishment of a functional barrier to macromolecules and pathogenic bacteria.
15 This phenomenon is called gut closure and appears to be affected by the diet. Hence, different studies with infants (JPGN, 1995, 21: 383-6) and animal models (Pediatr Res, 1990, 28: 31-7) show that the maturation of the barrier is faster in breast-fed than in formula-fed newborns. This could explain the higher prevalence of allergy and infection in infants fed formula than in those fed with mother milk.

20 An impressive number of different mechanisms integrates this barrier, mechanisms that act synergically to protect the host from the luminal aggressions. The first barrier consists on the intestinal epithelium, a continuous monolayer of columnar epithelial cells sealed together by protein complexes, such as the tight junctions. The second is a non-specific barrier composed by
25 mechanisms that protect the mucosal surface as saliva, gastric acidity, mucus layer, proteolytic digestion, alkaline intestinal pH, unstirred layer and intestinal peristalsis. The gut immune system (GALT) is able to respond selectively and specifically to the foreign molecules and pathogen microorganisms. Finally, and not less important, intestinal flora directly and indirectly protect against host invasion by pathogens and macromolecules with antigenic properties.

30 Moreover, physical stress due for instance to antibiotherapy, disease or surgery and psychological stress induced for instance by hospitalization or prolonged separation from the mother, common situations in the preterm infant population, may impair further the maturation of the intestinal

NO 7570/EP

barrier and delay closure. Therefore, microbial flora has shown to affect the status of various mechanisms of the intestinal barrier (mucus layer, tight junctions) and both physical and psychological stress have been related to increased permeability to macromolecules, small solutes and bacteria.

5

In the art several means have been proposed to improve gut barrier function or gastrointestinal health in infants. For example, in US 6132710, purified *Lactobacillus salivarius* and *Lactobacillus plantarum* strains are administered to preterm infants to prevent injury caused by infection and inflammation to mucosal tissue, especially by nasogastric administration to prevent
10 gastrointestinal tissue injury in neonatal necrotizing enterocolitis.

15

Also, JP 5030942 provides a food and drink containing active component of milk fat globule membrane (MFGM), which can control permeability of high molecular substances, such as protein, through intestine vessel. It is useful for prevention and therapy of food allergic disease.

20

As regards to gut barrier immunity, WO 9700078 provides a protein hydrolysate for down-regulating hypersensitivity reactions and for promoting the gut immune barrier, which is made by hydrolyzing proteins with: (a) enzymes derived from a preparation containing probiotic gastrointestinal bacteria and (ii) a protease system similar to that of *Lactobacillus GG* and (b)
20 pepsin and/or trypsin.

25

Though those microorganisms or ingredients bring about a great potential of beneficial effects for the individual incorporating them, a disadvantage resides in that said microorganisms or ingredients only exert their effect in limited parts of the intestine and on individual mechanisms
25 of the intestinal barrier.

30

Therefore, an object of the present invention resides in obviating the disadvantages of the prior art and providing improved means to promote simultaneous maturation of various mechanisms of the gut barrier and this all along the intestine during formula feeding, in order to induce a
30 pattern of gut barrier maturation similar to that observed with breast-feeding.

Summary of the invention

NO 7570/EP

During the studies leading to the present invention the present inventors have realized that the above object may be solved by providing specific combinations of bioactive ingredients that can be associated with microorganisms able to deliver at least one of said ingredients all along the intestine.

These specific combinations of ingredients comprise at least one substance selected from the group consisting of milk fractions, protein hydrolyzates, specific fats (gangliosides, LC-PUFAs), polyamines, non-digestible oligosaccharides.

10

The specific microorganisms according to the present invention that may be used as part of the combinations are not meant to survive in the gastro-intestinal tract and colonize it, but rather the present invention proposes the use of microorganisms for delivering said substances able to improve the gut barrier maturation to specific parts of the gastro-intestinal tract thereof. The microorganisms differing in their ability to survive in the different parts of the gastro-intestinal tract can be incorporated in a cocktail. Some of said substances can be added to the microorganism cocktail in order to reinforce their effects by stimulating the maturation of barrier mechanisms different to those stimulated by the microorganisms.

These combinations of ingredients could be included in pre-term and starter infant formulas that, by improving barrier maturation could also reduce the risk of allergy and infection.

Thus, according to the present invention, a nutritional formula designed for improving gut barrier maturation and an optimal barrier function in infants, comprises one of the above combinations of ingredients, supplemented in an amount efficient to induce a pattern of gut barrier maturation similar to that observed with breast-feeding.

A further object of the present invention relates to the use of such specific ingredient combinations to improve gut barrier maturation and barrier function in infants, favorable to infant health and thus reducing the risk of developing allergy and infection.

In another embodiment, the present invention relates to the use of the said ingredients to improve the gut barrier homeostasis during the suckling period in healthy infants and in those suffering from physical or psychological stress.

- 5 In a last aspect, the invention provides a method for improving gut barrier maturation and an optimal barrier function in infants, comprising the step of administering to the individual a combination of at least one ingredient selected from the group consisting of milk fractions, protein hydrolyzates, specific fats, polyamines, non digestible oligosaccharides said ingredients being associated with microorganism as delivering agent where necessary.

10

Detailed description of the invention

- In the following description, the term "Low birth weight (LBW) formula" means formula specifically designed for feeding the LBW infant. The LBW infant is defined as an infant
15 weighing less than 2500 g at birth. This infant may be either a "premature" infant (i.e. born before the 37th week of gestation) or a "small-for-date" infant (i.e. an infant born between the 37th and 41st week of gestation but showing a retarded intra-uterine growth). The LBW formula can be used as soon as enteral feeding is possible and until the LBW infant achieves a body weight similar to the birth weight of full-term infant (2500 g- 4000 g) or for the further weeks
20 until 5000 g.

The term "Starter formula" means formula specifically designed for feeding infants during the first 4-6 months of life and fulfilling the totality of their nutritional requirements.

- 25 According to a first aspect, the following substances may be part of the combination which can improve barrier maturation all along the intestine during formula feeding :
- Non-digestible carbohydrates, such as fructo-oligosaccharides (FOS), galacto-oligosaccharides (GOS), inulin, Arabic gum, resistant starch and the like
 - Human milk oligosaccharides, such as sialyllactose and/or
 - 30 - LC-PUFA, such as arachidonic acid (AA) or docosahexanoic acid (DHA) and/or
 - Gangliosides such as those contained in delactosed whey from buffalo milk, and/or

NO 7570/EP

- Milk or colostrum fractions, such as acid, rennet or micellar casein, acid, sweet or ultra whey, fat globules membranes and the like, and/or
- Extensive hydrolysed protein, such as those obtained from whey protein hydrolysis, and/or
- 5 - Polyamines such as spermine or spermidine and/or
- Microorganisms.

Preferably, non-digestible carbohydrates may be selected in the group of fructo-oligosaccharides, galacto-oligosaccharides, sialo-oligosaccharides, xyloligosaccharides, inulin, arabic gum, guar gum, resistant starch and/or milk-derived oligosaccharides and be added to the microorganism cocktail. One or more of these can be used in the total doses of from about 0.01 to 5 g/100 ml, and preferably 1-2 g/100 ml. A mixture of two or more carbohydrates may be used, each carbohydrate ranging between the 5% to 95% of the carbohydrate mixture.

15 Preferably, particular lipids may be used. For instance, an effective amount of at least one n-6 polyunsaturated fatty acid in combination with at least one n-3 polyunsaturated fatty acid, such as C20 or C22 n-6 fatty acid and one C20 or C22 n-3 fatty acid. The C20 or C22, n-6 fatty acid is present in a total amount of about 0.01 to 6.0% by weight of all fatty acids in the composition, preferably in a total amount of 0.1 to 1%. The C20 or C22 n-3 fatty acid is included in a total amount of about 0.01 to about 6.0% by weight of all fatty acids in the composition, preferably in a total amount of 0.1 to 1%. Preferably, the n-6 polyunsaturated fatty acid used in the present invention is arachidonic acid (AA, C20:4 n-6) and the n-3 polyunsaturated fatty acid used in the present invention is docosahexanoic acid (DHA, C22:6, n-3). The effective AA: DHA ratio is about 1: 1 to 2.5: 1, and preferably 1: 1 to 2: 1. The source of the LC-PUFA may be egg lipids, 25 fungal oil, low EPA fish oil, algal oil, etc.

Gangliosides, a second class of lipids, may also be added to the combination of ingredients, for example in an amount of from about 1-20 microMol/L formula, and preferably 6-15 microMol/L. The source of gangliosides may be cows milk, cows colostrum, but preferably buffalo milk, 30 buffalo milk serum, buffalo colostrum, buffalo colostrum serum and/or derivatives of either.

The combination may also contain polyamines, in particular spermidine, spermine, or putrescine

NO 7570/EP

and/or one or more polyamine precursors, in particular ornithin and arginine. They can be used in an amount of about 10 to 2,000 microg/100 g solid formula. The polyamine is preferably at least two or more selected from the group consisting of spermine, spermidine, putrescine and cadaverine. Preferably the composition comprises about 10-90% of spermine, 10-90% of spermidine, 0-90% of putrescine and 0-20% of cadaverine.

Preferably, the milk fractions (enriched in growth factors) may be in the form of fat globule membrane proteins, acid, rennet or micellar casein, acid, sweet or ultra whey, whey protein hydrolysates, for example. They can be used in an amount of about 0.01 to 7 g /100 ml formula, and preferably 0.5-3 g/100 ml.

According to another aspect, any or several of the former substances may be associated with microorganisms as delivering agents.

The micro-organisms to be used contain at least one substance the release thereof at the specific location will result in a beneficial effect on the barrier maturation. The microorganisms to be used can be specifically designed, treated or modified to ensure the release at the specific location.

Examples for a specific delivery to the small intestine are e.g. substances that interact locally with the mucus layer of the host, aggregate pathogens and facilitate their elimination by mucus flushing substances, e.g. substances that complex macromolecules and reduce their ability to permeate, e.g. enzymes that have the property to digest pathogen virulence factors (such as enterotoxins). Examples for a delivery to the colon are e.g. substances that have detoxifying properties, substances that have the potential to control the motility pattern of specific gut portions, substances that have the potential to favor intestinal cell differentiation, such as polyamines, substances that have the potential to increase innate immunity or substances that have the potential to restore the mucus layer integrity.

In order to provide a microorganism containing one or more substances of interest any microorganism may be selected, that inherently expresses such substances. Since the microorganisms are designed to release their intracellular material including the beneficial substance(s) at a specific location of the gut, a secretion of the substance into the environment is

NO 7570/EP

not required. On the contrary, according to the present invention the substance will be present in higher amounts at the predetermined location, since essentially all of the microorganism utilized will lyse and release the substance there. To this end, the corresponding microorganisms already containing the respective substance may optionally be pretreated in a manner appropriate to deliver the substance up to a certain desired location of the gut and may be administered to a recipient, whereupon they will lyse at the respective location in the gut depending on the sort of pretreatment.

This is a great advantage as compared to the common use of probiotics, wherein beneficial substances are primarily released by means of secretion into the environment. According to the present invention the microorganism utilized will release all of its beneficial cargo essentially at the same time when arriving at the location of the gut, where it is designed to lyse. In addition, the amount of the corresponding substance to be delivered to a recipient may also be more properly controlled, since a given amount of the microorganism to be used will be administered, with the content of the substance of interest being by and large known.

In order to increase the amount of the said substances to be delivered by the microorganism common techniques may be used, such as: applying particular fermentative conditions or genetically modifying the microorganism itself, by e.g. subjecting the microorganisms to a random mutagenesis and selecting those mutants expressing a higher amount of the desired substance. Yet, also recombinant means may be applied, wherein the expression of the endogenous gene is increased by e.g. linking the corresponding gene with a promotor stronger than the endogenous one, or by inserting the gene or genes encoding the substance(s) of interest into the microorganism on a plasmid or into the chromosome thereof, optionally linked with a strong promoter that drives the expression of the gene(s) of interest such that the recombinant microorganism will contain higher amounts of the desired substance.

Depending on the nature and duration of the pretreatment the endurance of the microorganism, i.e. its survival in the gastro-intestinal tract may be established, with potential locations of delivery being the stomach, the duodenum, the jejunum, the ileum or the colon. The microorganisms to be added in the present formula may be selected from the group consisting of Lactobacilli, Bifidobacteria, Streptococci, Pediococci, Enterococci, Lactococci, Oenococci,

NO 7570/EP

Staphylococci, Bacteroides, or mixtures thereof. Preferred examples of such microorganisms are Bad 4, B128, B129 or *Lactobacillus johnsonii* all of which are freely available from Depository Institutes under the accession nos. CNCM I-2168, CNCM I-2169, CNCM I-2170 or CNCM I-1225 respectively. Also, *Streptococcus thermophilus* (TH4) or *Bifidobacterium lactis* (Bb12 (ATCC27536)) may be used. They are provided by Hansen (Chr. Hansen A/S, 10-12 Boegs Alle, P.O. Box 407, DK-2970 Hoersholm, Denmark).

Once a microorganism has been selected and optionally pretreated, said microorganism may be included in a LBW or starter formula as a powder obtained by freeze- or spray-drying, for example in an amount of from 10^5 - 10^{13} cfu/100 g, depending on the nature of the substance to be delivered and the amount of the substance contained in the respective microorganisms.

The above ingredients are conveniently administered in form of a product acceptable to the consumer, such as an ingestible carrier or support, respectively. Examples for such carriers or supports are a pharmaceutical or a food or petfood composition. Non-limiting examples for such compositions are milk, yogurt, curd, cheese, fermented milks, milk based fermented products, fermented cereal based products, milk based powders, infant formula, liquid bacterial suspensions, dried oral supplement, wet oral supplement, dry tube feeding or wet tube feeding.

The nutritional compositions are preferably in the form of a complete diet such that, when used as the sole source of nutrition, essentially covers all daily energy, nitrogen, lipid, vitamin, mineral and trace elements. However, the nutritional composition may also be in the form of a supplement.

In a preferred embodiment, the present invention provides an infant formula, which may be in the form of a low birth weight or a starter infant formula, for example. It may comprise apart the combination of specific ingredients as mentioned above, a protein source, a carbohydrate source, and a source of lipids.

The source of protein may be any suitable dietary protein; for example animal proteins (such as milk proteins, meat proteins and egg proteins), vegetable proteins (such as soy, wheat, rice or pea proteins), mixtures of free amino acids, or combination thereof. Milk proteins such as casein,

NO 7570/BP

whey proteins and soy proteins are particularly preferred. In a preferred embodiment, the protein source comprises about 1.8 to about 4 grams per 100 kcal of formula.

If the formula includes a fat source, the fat source preferably provides about 5% to about 55% of the energy of the nutritional formula; or about 3 to 7 grams per 100 kcal of formula; The lipids making up the fat source may be any suitable fat or fat mixture. Vegetable fats are particularly suitable; for example soy oil, palm oil, coconut oil, safflower oil, sunflower oil, corn oil, canola oil, lecithins, and the like. Animal fats such as milk fats may also be added if desired.

If the formula includes a carbohydrate source, the carbohydrate source preferably provides about 40% to about 80% of the energy of the nutritional formula or about 6 grams to about 15 grams per 100 kcal of formula, for example. Any suitable carbohydrates may be used, for example sucrose, lactose, glucose, fructose, corn syrup solids, and maltodextrins, and mixtures thereof.

Suitable vitamins and minerals may be included in the nutritional formula in the usual manner to meet the appropriate guidelines. One or more food grade emulsifiers may be incorporated into the nutritional formula if desired; for example diacetyl-tartaric acid esters of mono-diglycerides, lecithin and mono- and di-glycerides. Similarly suitable salts and stabilisers may be included.

This formula is preferably enterally administrable; for example in the form of a powder, a liquid concentrate, or a ready-to-drink beverage. It may be prepared in any suitable manner, for example, by blending together the source of dietary protein, the carbohydrate source, and the fat source in appropriate proportions. If used, the emulsifiers may be included in the blend. The vitamins and minerals may be added at this point but are usually added later to avoid thermal degradation. Any lipophilic vitamins, emulsifiers and the like may be dissolved into the fat source prior to blending. Water, preferably water which has been subjected to reverse osmosis, may then be mixed in to form a liquid mixture. The temperature of the water is conveniently about 50°C to about 80°C to aid dispersal of the ingredients. Commercially available liquefiers may be used to form the liquid mixture. The liquid mixture is then homogenized; for example in two stages.

The liquid mixture may then be thermally treated to reduce bacterial loads. For example, the liquid mixture may be rapidly heated to a temperature in the range of about 80°C to about 150°C for about 5 seconds to about 5 minutes. This may be carried out by steam injection, autoclave or by heat exchanger; for example a plate heat exchanger. The liquid mixture may then be cooled to

NO 7570/EP

about 60°C to about 85°C; for example by flash cooling. The liquid mixture may then be again homogenized; for example in two stages at about 7 MPa to about 40 MPa in the first stage and about 2 MPa to about 14 MPa in the second stage. The homogenized mixture may then be further cooled to add any heat sensitive components; such as vitamins and minerals. The pH and solids
5 content of the homogenized mixture is conveniently standardized at this point.

If it is desired to produce a powdered nutritional formula, the homogenized mixture is transferred to a suitable drying apparatus such as a spray drier or freeze drier and converted to powder. The powder should have a moisture content of less than about 5% by weight.

- 10 If it is desired to produce a liquid formula, the homogenized mixture is preferably aseptically filled into suitable containers. Aseptic filling of the containers may be carried out by pre-heating the homogenized mixture (for example to about 75 to 85°C) and then injecting steam into the homogenized mixture to raise the temperature to about 140 to 160°C; for example at about 150°C. The homogenized mixture may then be cooled, for example by flash cooling, to a
15 temperature of about 75 to 85°C. The homogenized mixture may then be homogenized, further cooled to about room temperature and filled into containers. Suitable apparatus for carrying out aseptic filling of this nature is commercially available. The liquid formula may be in the form of a ready to feed formula having a solids content of about 10 to about 14% by weight or may be in the form of a concentrate; usually of solids content of about 20 to about 26% by weight. Flavors
20 may be added to the liquid formulas so that the formulas are provided in the form of convenient, flavorful, ready-to-drink beverages.

- This composition may be particularly designed for healthy infants, infants suffering from gut microflora alterations, such as after antibiotic treatment and, infants suffering from physical and
25 psychological stress resulting, for example, from disease, surgery, hospitalization, prolonged separation from the mother, in order to improve gut barrier maturation and thus reduce the risk of allergy and infection. The amount of the formula required to be fed to the infant will vary depending upon factors such as the infant's condition, the infant's body weight, the age of the infant, and whether the formula is the sole source of nutrition. In general, sufficient of the
30 nutritional composition is administered to provide the infant with about 1 g protein to about 4.0 g protein per kg of body weight per day supplemented with the ingredients according to the present invention in the amounts as indicated above. If the nutritional composition is used as a

NO 7570/EP

supplement to other foods, the amount of the nutritional composition that is administered daily may be decreased accordingly.

The following non-limiting examples further illustrate the invention.

5 Example 1: Formula for low-birth-weight infants

The formula has the following composition (per 100 g of powder) : total fat 24g, total protein 14.4 g, total carbohydrates 55.9 g, AA enriched oil (fungal) 0.87g, DHA enriched oil (Low EPA fish oil) 0.44g, FOS/inulin (70/30) 12g, S. thermophilus (freeze-dry powder, 10E12cfu/g) 0.1g,
10 B. lactis (freeze-dry powder, 5x10E12cfu/g) 0.15g, Spermine/Spermidine mix (1/1) 0.1mg, Sodium 180 mg, Potassium 530 mg, Chloride 280 mg, Phosphorus 320 mg, Calcium 490 mg, Magnesium 54 mg, Manganese 34 µg, Vitamin A 1500 IU, Vitamin D 490 IU, Vitamin E 9.8 IU, Vitamin C 79 mg, Vitamin K1 59 µg, Vitamin B1 0.29 mg, Vitamin B2 0.66 mg, Vitamin B6 0.37 mg, Niacin 4.9 mg, Folic acid 290 µg, Pantothenic acid 2.3 mg, Vitamin B12 1.1 µg, Biotin
15 11 µg, Choline 37 mg, Inositol 22 mg, Taurine 39 mg, Carnitine 7.9 mg, Iron 7.4 mg, Iodine 49 µg, Copper 0.44 mg and Zinc 3.7 mg.

The formula is reconstituted by mixing 142 g of powder to 900 mL of water to give 1 L of ready-to-drink preparation. The composition given above can vary to accommodate for local directives
20 concerning the amounts of specific ingredients. Other trace elements (e.g. selenium, chromium, molybdenum, fluoride) may be added in adequate amount according to age.

Example 2 : starter formula

A starter formula for infants (from birth to 4-5 months), in powder form is prepared. The formula
25 has the following composition (per 100 g of powder): total fat 25.8 g; total protein 11.5 g; total carbohydrates 57.8 g, AA enriched oil (fungal) 1g, DHA enriched oil (Low EPA fish oil) 1g, FOS/inulin (70/30) 12g, L. paracasei (Spray-dry powder, 10E12cfu/g) 0.1g, B. longum (Spray-dry powder, 5x10E12cfu/g) 0.1g, Sodium 120 mg, Potassium 460 mg, Chloride 360 mg, Phosphorus 160 mg, Calcium 320 mg, Magnesium 35 mg, Manganese 40 µg, Vitamin A 1500 IU, Vitamin D
30 310 IU, Vitamin E 6.1 IU, Vitamin C 41 mg, Vitamin K1 42 µg, Vitamin B1 0.31 mg, Vitamin B2 0.69 mg, Vitamin B6 0.38 mg, Niacin 3.8 mg, Folic acid 46 µg, Pantothenic acid 2.3 mg,

Vitamin B12 1.1 µg, Biotin 11 µg, Choline 38 mg, Inositol 23 mg, Taurine 41 mg, Carnitine 8.2 mg, Iron 6.1 mg, Iodine 25 µg, Copper 0.31 mg and Zinc 3.8 mg,

5 The formula is reconstituted by mixing 132 g of powder to 900 mL of water to give 1 L of ready-to-drink preparation. The composition given above can vary to accommodate for local directives concerning the amounts of specific ingredients. Other trace elements (e.g. selenium, chromium, molybdenum, fluoride) may be added in adequate amount according to age.

Example 3: starter infant formula

10

A starter formula for infants is prepared as in example 2, but replacing FOS/inulin by sialyl-lactose in an amount of 0.5 g. In this formula, half of the total protein will be furnished in the form of extensive whey protein hydrolyzate.

15

Claims

1. A nutritional composition intended for improving gut barrier maturation and ensuring an optimal barrier function in infants, comprises a combination of a least one substance
5 selected from the group consisting of milk fractions, protein hydrolyzates, specific fats, polyamines or non digestible oligosaccharides, in an amount efficient to induce a pattern of gut barrier maturation similar to that observed with breast-feeding.
2. A composition according to claim 1, wherein at least one of said substances is associated
10 with microorganism as delivering agent, in particular *Lactobacilli*, *Bifidobacteria*, *Streptococci*, *Pediococci*, *Enterococci*, *Lactococci*, *Oenococci*, *Staphylococci*, *Bacteroides*, or mixtures thereof.
3. A composition according to claim 2, wherein the microorganism is *Bifidobacterium*
15 *CNCM I-2170*, *Bifidobacterium CNCM I-2168*, *Bifidobacterium CNCM I-2169*, *Lactobacillus johnsonii CNCM I-1225*, *Lactobacillus paracasei CNCM I-2116*, *Bifidobacterium lactis ATCC 27536*, *Streptococcus thermophilus* (Chr. Hansen) or mixtures thereof.
- 20 4. A composition according to any of preceding claims, wherein the composition is in the form of a product acceptable to the consumer, such as an ingestible carrier or support, such carriers or supports being a pharmaceutical or a food or petfood composition.
- 25 5. A composition according to any preceding claims, wherein the composition is in the form of a complete diet, a supplement or a medicament.
6. A composition according to any preceding claims, wherein the composition is a low birth weight or a starter infant formula.
- 30 7. Use of a combination of a least one substance selected from the group consisting of milk fractions, protein hydrolyzates, specific fats, polyamines or non digestible

oligosaccharides, for the preparation of a composition intended for improving gut barrier maturation and barrier function, particularly in infants.

- 5 8. Use of a combination of a least one substance selected from the group consisting of milk fractions, protein hydrolyzates, specific fats, polyamines or non digestible oligosaccharides, for the preparation of a composition intended for reducing the risk of developing allergy and infection.
- 10 9. Use of a combination of a least one substance selected from the group consisting of milk fractions, protein hydrolyzates, specific fats, polyamines or non digestible oligosaccharides, for the preparation of a composition intended for improving the gut barrier homeostasis during the suckling period.
- 15 10. The use according to any of preceding claims, wherein at least one of said substances is associated with microorganism as delivering agent, in particular *Lactobacilli*, *Bifidobacteria*, *Streptococci*, *Pediococci*, *Enterococci*, *Lactococci*, *Oenococci*, *Staphylococci*, *Bacteroides*, or mixtures thereof.
- 20 11. The use according to any of preceding claims, in which the microorganism is *Bifidobacterium* CNCM I-2170, *Bifidobacterium* CNCM I-2168, *Bifidobacterium* CNCM I-2169 *Lactobacillus johnsonii* CNCM I-1225, *Lactobacillus paracasei* CNCM I-2116, *Bifidobacterium lactis* ATCC 27536, *Streptococcus thermophilus* (Chr. Hansen) or mixtures thereof.
- 25 12. A method for improving gut barrier maturation and an optimal barrier function in infants, comprising the step of administering to the individual a combination of at least one ingredient selected from the group consisting of milk fractions, protein hydrolyzates, specific fats, polyamines, non digestible oligosaccharides in an amount efficient to induce a pattern of gut barrier maturation similar to that observed with breast-feeding.
- 30 13. A method according to claim 10, wherein at least one of said substances is associated with microorganism as delivering agent.

Abstract**Nutritional formula for optimal gut barrier function**

- 5 The present invention pertains to a composition for inducing a pattern of gut barrier maturation similar to that observed with breast-feeding and able to improve gut barrier maturation, e.g. during neonatal stress. In particular, the present invention relates to an infant formula containing a combination of specific ingredients designed to provide a synergistic effect all along gastrointestinal tract and barrier function.

This Page Blank (uspto,

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

This Page Blank (uspto)